

4° CONGRESSO NAZIONALE FRAGILITY FRACTURE NETWORK - ITALIA

Appropriatezza, Qualità e Sostenibilità delle Cure nel Percorso Ortogeriatrico



LUNCH SYMPOSIUM

Salvaguardare la relazione pazientecaregiver

Dott. Stefano Eleuteri

Health Psychologist, PhD in Education FFN EduCom Chair, FFN Italy EduCom Chair Lecturer













INTRODUCTION





- Fragility fracture is one of the major causes of loss of self-sufficiency in older patients;
- Most of patients hospitalised with hip fracture are discharged quickly after surgery and therefore the rehabilitation task of caregiving often "falls on the shoulders" of a caregiver within the family.





THE CAREGIVERS





As a result of reduced ability to complete activities of daily living, individuals who have suffered a fragility fracture may rely on informal caregivers

During the first year after a fracture, the hours of care provided by relatives vary greatly by fracture type: the more serious the fracture, the more support is needed

(Svebdom et al., 2018; figure taken from IOF Report, 2018)




CULTURE INFLUENCE



The hours of care provided by relatives vary greatly also by country


In countries where crossgenerational support is more established, the impact of fragility fractures on caregivers is generally higher




(EUROCARERS, 2018; figure Modified from IOF Report, 2018)



Psychological wellbeing in HF patients and their carers






(Eleuteri et al., 2016)


Lower levels of depressive disorders and of anxiety disorders and higher levels of health, vitality, positivity and self-control in the caregiver correspond to a better functional outcome in the patient (p<0.01)




Psychological wellbeing in HF patients and their carers



The CBI total score is negatively related with patient's PGWBI score






Patient's PGWBI score is related with the caregiver's PGWBI General Health subscale score

(Eleuteri et al., 2016)



Mobility capacity



Differences in ADL score and autonomus mobility capacity in participants whose caregivers reported higher or lower burden

	ADL2	WALKING
Mann-Whitney U	209,00	256,00
Wilcoxon W	485,00	556,00
Z	-2,618	-2,273
Asymp. Sig. (2-tailed)	,009**	,023*


(Eleuteri et al., 2016)



Health education



- We also have evidence that health education can be effective in improving functional recovery and patient adherence.
 Has been identified as a key element of the effectiveness of fracture coordination units.
- Health education aimed at empowering the patient and caregiver for a safe transition to home, together with the support of these home hospitalisation units that could provide support during the first days of discharge home, could contribute to a reduction in referrals and an improvement in the functional recovery and quality of life Rojas-Ocaña et al., 2021; Rojas-Sánchez et al., 2009; Guevara et al., 2011; Azzolin et al., 2013; Corrales & y Sánchez, 2018; Hernández Bernal et al., 2018



SUPPORTING FAMILY CAREGIVERS OF OLDER ADULTS WITH HIP FRACTURE: THE ROLE OF EDUCATIONAL COURSES

Maria Eduarda Batista de Lima¹, Stefano Eleuteri² and Paolo Falaschi³

¹ Fragility Fracture Network, Education Committee Nurse Member
² Department of Psycology, Sapienza, University of Rome, Italy; Fragility Fracture Network, Education Committee Deputy Chair
³ Fragility Fracture Network, Past President and Education Committee Chair

urnitin


Description Reviews



Traduzione e adattamento: Stefano Eleuteri

Al termine di questo modulo il caregiver sarà in grado di:

- Definire il termine "caregiver"
- Conoscere i sintomi correlati al sovraccarico assistenziale del caregiver
- Comprendere la connessione tra la salute psicologica della persona assistita e quella del caregiver
- Conoscere le strategie per prevenire il sovraccarico assistenziale e per essere un caregiver competente
- Riconoscere l'importanza delle reti di supporto nel processo di assistenza



Traduzione e adattamento: Stefano Eleuteri

Description

Al termine del modulo, il caregiver sarà in grado di:

- · Definire il termine "stress":
- Conoscere i sintomi e le cause dello stress del caregiver;
- · Sapere come reagire allo stress;
- Comprendere e conoscere le strategie per la gestione dello stress.

Reviews





■ Corso digitale per una cura consapevole dei pazienti con fratture da fragilità





Survey rivolta ai medici specialisti e ai medici in formazione per indagare la gestione e le caratteristich cliniche dei pazienti anziani con cardiopatia ischemica cronica.

Link: https://redcap.link/SIGG_CIC

sigg.it/news/survey-sigg/

Gruppo di Studio sulla prevenzione del maltrattamento nell'anziano VALUTAZIONE DEL BURNOUT DEL PERSONALE E COMPORTAMENTI POTENZIALMENTE INADEGUATI VERSO GLI ANZIANI ACCOLTI IN RSA

Il questionario è anonimo e può partecipare il personale sanitario ed assistenziale (medici, infermier operatori socio sanitari, fisioterapisti, educatori, psicologi, altri operatori socio-assistenziali)

Il questionario è accessibile al SEGUENTE LINK

4° CONGRESSO NAZIONALE FRAGILITY F





cost-programming.eu/survey/

>20 languages

Retweet in your native language!
#COST #COSTactions #ScienceWithoutBorders
@COSTprogramme





